			BEST A	AVAILABLE CO	PY	10	/049822
Sheet 1sof 1					JU 13 Re	通 程[[1]]	1 9 FEB 2002
FORM PTO 1449 (modified) U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE LIST OF REFERENCES CITED BY APPLICANT(S)				ATTY DOCKET NO. 2002_0256A NEW			
				APPLICANT Shigeo OHTA et al.			
(Use several sheets if necessary) Date Submitted to PTO: February 19, 2002				FILING DATE February 19, 2002		GROUP	
				U.S. PATENT DOCUMENTS			
*EXAMINER INITIAL		DOCUMENT NUMBER	DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
	AA						
	AB						
	AC						
	AD		<u></u>				
	-			FORFIGN PATENT DOCUMENTS			
	ļ	DOCUMENT NUMBER	DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION YES NO
-	AE						
	AF						
		OTH	ER DOCUMENTIS	6) (Including Author, Title, Date, Po	ertinent Pages, Etc.)		
/CY/	AG	1997.					
	АН						
	AI	D.A.M. Grillot et al., "Genomic Organization, Promoter Region Analysis, and Chromosome Localization of t Mouse bc1-x Gene", J. Immunol., Vol. 158, No. 10, pp. 4750-4757, 1994.					
	LA	N. Shirakawa et al., "An Additional Form of Rat Bc1-x, Bc1-x β, Generated by an Unspliced RNA, Promotes Apoptosis in Promyeloid Cells", J. Biol. Chem., Vol. 271, No. 22, pp. 13258-13265, 1996.					
\downarrow	AK	M. Gonzalez-Garcia et al., "bc1-x is the Major bc1-x mRNA form Expressed During Murine Development and its Product Localizes to Mitochondria", Vol. 120, Development, No. 10, pp. 3033-3042, 1994.					
/CY/	AL	J. L. Tilly et al., "Expression of Members of the Bc1-2 Gene Family in the Immature Rat Ovary: Equine Chorionic Gonatropin-Mediated Inhibition of Granulosa Cell Apoptosis is Associated with Decreased Bax and Constitutive Bc1-2 and Bc1-x _{kmg} Messenger Ribonucleic Acid Levels", Endocrinology, Vol. 136, No. 1, pp.					

/Christopher Yaen/ (04/02/2007)

04/02/2007